



Williams, E. J., & Joinson, A. (2020). Developing a Measure of Information Seeking about Phishing. *Journal of Cybersecurity*, 6(1), [tyaa001]. <https://doi.org/10.1093/cybsec/tyaa001>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1093/cybsec/tyaa001](https://doi.org/10.1093/cybsec/tyaa001)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Oxford University Press at <https://doi.org/10.1093/cybsec/tyaa001> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>


A 60km-resolution GCM for Large-Ensemble Climate Simulations in *climateprediction.net*

Peter Watson (Oxford University)

with William Ingram, Sarah Sparrow, Simon
Wilson, Marie Drouard, Giuseppe Zappa,
Richard Jones, Daniel Mitchell, Tim
Woollings, Myles Allen

The *climateprediction.net* infrastructure


- At present, the Met Office models HadXM3(P) are run on thousands of volunteers' computers using distributed computing to study extreme weather events.
- Used because they are highly memory-efficient.
- The global model has ~150km L19 resolution and regional models have 25km or 50km resolutions.



The need for higher global resolution


- Many extreme weather events involve large-scale dynamical extremes e.g. blocks, storms.

High blocking occurrence in Europe in 2003 summer:



Mitchell et al. (2016)

High number of storms in 2013/14 winter:



- Simulation of these events improves considerably going from ~100km to ~20-40km global resolution (e.g. Jung et al., 2012; Davini et al., 2017)

DOCILE project


- Aim is to develop ~60km (N216) and ~90km (N144) 38-level global atmospheric models to run in *climateprediction.net*.
- Will be open to use by the academic community.
- Based on Met Office's HadAM4 (Williams et al., 2003).
- HadAM4 also includes updates to parameterisation schemes (microphysics, clouds and boundary layer).

Role for statistics

- Model evaluation
- Finding optimal parameters
- Getting information from the multi-thousand member ensembles.

DJF SAT and Precip biases

- Comparing atmosphere-only HadAM4 60km (3 members) and state-of-the-art HadGEM3 GA6 60km (15 members; Ciavarella et al. [2018]) biases over 1986-2012.
- Using GPCP precip, CERES for radiation and ERA-Interim for other variables.






DJF U850 and U200 biases




HadAM4 60km

HadGEM3 60km

U850







U200




DJF dynamical variability

- Blocking bias
 - HadAM4 shows the common negative bias over Europe, but it is comparable with HadGEM3's
 - Smaller than in the CMIP5 multi-model mean (Mitchell et al., 2016).
- Storm track density
 - HadAM4 captures main features well.
 - Biases have similar size to those in HadGEM3 except in continental North America.




DJF N. Atlantic jet latitude distribution

- The trimodality of the frequency distribution of the latitude of the North Atlantic eddy-driven jet is captured in HadAM4 (unlike in most CMIP5 models).




JJA biases

- Large biases in NH land JJA SAT, precip and radiation with initial parameter settings.
- Seems associated with having too little cloud.




Tuning the mean state

- Using “algorithmically-assisted” method (Neelin et al., 2010).
- Based on optimising a cost function that depends on RMSE of mean climate.
- 26 parameters perturbed – only the 8 most important were optimised.




Neelin et al. (2010)



Bellprat et al. (2016)

Preliminary tuning results

- Optimising 30-90N SAT, precip, U200 and U850 in JJA and DJF gives good results (90km resolution).



Thoughts on tuning methods

- Perturbing individual parameters alone helpful for debugging – useful to include as well as multi-parameter perturbations.
- Cost function that matches our best judgement is complex and uncertain (Hourdin et al., 2016) – counts in favour of using emulator (e.g. Rougier et al., 2009; Neelin et al., 2010) rather than iterative updating (e.g. Tett et al., 2013).
- Methods that take into account whole distributions would be useful.

Summary

- 60km and 90km L38 configurations of HadAM4 have been newly developed for the *climateprediction.net* distributed computing project.
- This will allow large-ensembles to be produced to study extreme events with a much better representation of extratropical dynamics.
- HadAM4 can simulate NH extratropical DJF weather about as well as a state-of-the-art Met Office model.
- Experiments to study NH winter extremes are in progress.
- Parameter tuning to reduce JJA model biases further is being finalised.